



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

Statistical description of glass network interconnectivity in calcium aluminosilicate glasses

Keding, Ralf; Moesgaard, Mette; Skibsted, Jørgen; Yue, Yuanzheng

Publication date:
2010

Document Version
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

Citation for published version (APA):
Keding, R., Moesgaard, M., Skibsted, J., & Yue, Y. (2010). *Statistical description of glass network interconnectivity in calcium aluminosilicate glasses*. Abstract from XXII International Congress on Glass, Bahia, Brazil.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

Statistical description of glass network interconnectivity in calcium aluminosilicate glasses

Ralf Keding, Mette Moesgaard, Jørgen Skibsted and Yuanzheng Yue

^aSection of Chemistry, Aalborg University, DK-9000 Aalborg, Denmark

^bDepartment of Chemistry, Aarhus University, DK-8000 Aarhus, Denmark

Two series of peralkaline glasses with a composition close to the eutectic compositions of anorthite-wollastonite-gehlenite and anorthite-wollastonite-tridymite were synthesized and investigated using ²⁹Si and ²⁷Al MAS-NMR. The structure of the glass is described by a network of SiO_{4/2} and AlO_{4/2}⁻ tetrahedrons as network former and Ca²⁺ ions as charge compensation for AlO_{4/2}⁻ or as network modifier. Besides the separation into bridging oxygen and non-bridging oxygen, the 2nd coordination sphere is described by a combinatorial statistic continuous random network or by a hierarchic random network. The description of the network is used to calculate the molar fraction of Si with different first (O) and second (Si or Al) coordination sphere. The ratio of the areas of the ²⁹Si NMR resonances of the different coordinated Si specimens is assumed to be proportional to the molar fraction obtained from the structural models. The other parameters for the peaks attributed to different ²⁹Si specimens are calculated by fitting the measured spectra. The quasi heterogeneous modeling approach with a hierarchy in the intermediate range order results in the best description of the NMR spectra while the intensities obtained from combinatorial statistic continuous random network do not result in a proper fit. This procedure of analyzing common ²⁹Si MAS-NMR gives hints, that the investigated Ca-aluminosilicate glasses are best described by a quasi-heterogeneous intermediate range order.